
# Fracture of the big toe

Hallux fracture





## **Table of contents**

Treatment	4
Instructions	4
Recovery	5
Reapply the pressure bandage	5
Exercises	5
Questions?	5

You have a fracture of the big toe (hallux). We treat this fracture with a pressure bandage and a special shoe.

#### Summary

- you have a fracture of the big toe
- this type of injury almost always heals well without any loss of function
- a routine follow-up is not necessary for this type of injury
- you can remove the pressure bandage if your pain allows you to do so,
   wear the pressure bandage for a maximum of 1 week
- to prevent the toe from getting stiff, it is important that you start moving the toe again as soon as possible
- you can find instructions on how to reapply the pressure bandage in the app
- if you have any questions, or are experiencing problems with your recovery, you can call the fracture hotline 'breuklijn' +3172 - 548 2465

#### Treatment

A fracture of the big toe (hallux) is a common injury. The fracture is treated with a pressure bandage and a special shoe that prevents you from putting weight on your toe.

week o - 1	pressure bandage and a special	
	shoe, remove as soon as pain	
	allows	
week 1 -3	• you may walk again wearing a	
	sturdy shoe	
after 3 weeks	• you may start sports, when your	
	pain allows you to do so	

#### Instructions

Follow these instructions for a good recovery:

- wear the pressure bandage for a maximum of 1 week. It is not a problem
  if you remove the pressure bandage earlier. If the pressure bandage gets
  loose, you can find instructions on how to reapply the pressure bandage in
  this app
- you may put pressure on your foot, if your pain allows you to do so. Use crutches if this is necessary
- elevate your foot when you are resting, for example, by putting it on a chair.
   This will reduce swelling and pain
- to prevent the toe from getting stiff, it is important that you start moving the toe again as soon as possible
- for the first three weeks, you should not practice any sports. After three
  weeks, you may start sports again if your pain allows you to do so. It can
  take up to 6-8 weeks before you have recovered enough to practice sports
  again
- are you in pain? Use a mild pain killer such as paracetamol (acetaminophen).
   Do not wait with the use of paracetamol until the pain is too much

#### Recovery

This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is unnecessary.

Does the pain increase, or does it not improve? Please contact the fracture hotline 'breuklijn': +31 72 - 548 2465.

#### ■ Reapply the pressure bandage

If you need to reapply the pressure bandage or shoe, you can find instructions on how to do so in in the Virtual Fracture Care app. You can buy a new pressure bandage at the pharmacy or reuse the pressure bandage that was applied in the Emergency Department.

#### Exercises

It is important to move the toe as much as possible. No specific exercises are necessary.

#### Questions?

#### You can call us

If there are any questions after reading the information, please contact us or visit <a href="https://www.nwz.nl">www.nwz.nl</a>

#### Contact Monday to Friday between 8.30-10.00 h and 13.30-15.00 h

• breuklijn: +31 72 548 2465

#### The Virtual Fracture Care app

Open the camera on your phone and aim it at the QR-code below. You will be automatically referred to the app.





Notes	5	

### Noordwest Ziekenhuisgroep

Postbus 501 1800 AM Alkmaar

www.nwz.nl

tel 072 - 548 4444

#### Colofon

Redactie chirurgie

communicatie

Druk Ricoh Bestelnummer 313045

Op alle onderzoeken en behandelingen van Noordwest Ziekenhuisgroep zijn de algemene voorwaarden van Noordwest Ziekenhuisgroep van toepassing, zie www.nwz.nl

