

# Collarbone fracture

Children

[www.nwz.nl](http://www.nwz.nl)



Noordwest  
Ziekenhuisgroep

# Table of contents

Treatment	4
Instructions	4
Recovery	5
Reapply the sling	5
Exercises	5
Questions?	5

**Your child has a fractured collarbone (clavicle). This fracture is treated with a sling.**

#### **Summary**

- your child has a fractured collarbone
- this type of injury almost always heals well without any loss of function
- a routine follow-up is not necessary for this type of injury
- you can remove the sling as soon as your child's pain allows you to do this. During the night you do not have to wear the sling. Use the sling for max. 4 weeks
- to prevent the shoulder from getting stiff, it is important that your child starts moving the shoulder and arm as soon as possible. Only use the shoulder when the pain allows this.
- you can find instructions on how to reapply the sling in the app
- if you have any questions, or are experiencing problems with your recovery, you can call the fracture hotline 'breuklijn':  
+31 72 - 548 2465

## ■ Treatment

A fractured collarbone is a common injury in children. The fracture is treated with a sling. If your child is young and won't wear the sling, this is not a problem. Collarbone fractures almost always heal without any problems or complications.

<b>week 0 - 4</b>	<ul style="list-style-type: none"><li>• wear the sling, remove it as soon as your child's pain allows this. Start moving the shoulder and arm as soon as possible, to prevent the shoulder from getting stiff. Only use the shoulder when the pain allows this</li></ul>
<b>week 4 - 6</b>	<ul style="list-style-type: none"><li>• let your child move the shoulder if the pain allows this</li></ul>
<b>after 6 weeks</b>	<ul style="list-style-type: none"><li>• gradually start sport activities</li></ul>

## ■ Instructions

Follow these instructions for a good recovery:

- the sling keeps your child's arm in a comfortable position for a few days to weeks
- you can remove the sling as soon as your child's pain allows you to do this. During the night you do not have to wear the sling. Use the sling for a maximum of 4 weeks
- to prevent a stiff shoulder it is important that your child starts moving his/her shoulder as soon as possible. Only use the shoulder when your child's pain allows you to do this
- activities such as sports can be done after 6 weeks when your child's pain allows you to do this. It is no problem to wait longer with these activities
- is your child in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until the pain is too much

## ■ Recovery

This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is unnecessary. As the bone heals, a bump may appear over the fracture. This is normal, it is the result of bone healing. This does not affect the function of your child's shoulder.

Does the pain increase, or does it not improve? Please contact the the fracture hotline 'breuklijn': +31 72 - 548 2465.

Physiotherapy is not necessary. If you are not satisfied with the function of your child's shoulder after 6 weeks, you can ask for a referral to a physiotherapist from your general practitioner. However, a referral is not required.

## ■ Reapply the sling

Instructions on how to reapply the sling can be found in the video in the Virtual Fracture Care app.

## ■ Exercises

To prevent a stiff shoulder it is important that your child starts moving the shoulder as soon as possible. Only use the shoulder when the pain allows you to do this. No specific exercises are necessary.

## ■ Questions?

### **You can call us**

If there are any questions after reading the information, please contact us or visit [www.nwz.nl](http://www.nwz.nl)

**Contact Monday to Friday between 8.30-10.00 h and 13.30-15.00 h**

- breuklijn: +31 72 - 548 2465

### The Virtual Fracture Care app

Open the camera on your phone and aim it at the QR-code below. You will be automatically referred to the app.



■ Notes

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

### Noordwest Ziekenhuisgroep

Postbus 501

1800 AM Alkmaar

[www.nwz.nl](http://www.nwz.nl)

**tel** 072 - 548 4444

### Colofon

Redactie            chirurgie  
                         communicatie

Druk                Ricoh

Bestelnummer    313061

*Op alle onderzoeken en behandelingen van Noordwest Ziekenhuisgroep zijn de algemene voorwaarden van Noordwest Ziekenhuisgroep van toepassing, zie [www.nwz.nl](http://www.nwz.nl)*